F44

NOAA Technical Report NMFS 44

September 1986

Synopsis of Biological Data on the Porgies, *Calamus arctifrons* and *C. proridens* (Pisces: Sparidae)

George H. Darcy

LIBRARY

OCT 21 1986

National Marine Fisheries Service
2570 Dole Street
Honolulu, Hawaii

FAO Fisheries
Synopsis No. 148

NMFS/S 148

SAST-*Calamus arctifrons* 1,70(39)034,02
C. proridens 1,70(39)034,06

NOAA
Tech.
Rept.
NMFS
44

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORT NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for their optimum use. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Reports series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of reports: Scientific investigations that document long-term continuing programs of NMFS; intensive scientific reports on studies of restricted scope; papers on applied fishery problems; technical reports of general interest intended to aid conservation and management; reports that review in considerable detail and at a high technical level certain broad areas of research; and technical papers originating in economics studies and from management investigations. Since this is a formal series, all submitted papers receive peer review and those accepted receive professional editing before publication.

Copies of NOAA Technical Reports NMFS are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from: U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161. Although the contents have not been copyrighted and may be reprinted entirely, reference to source is appreciated.

1. Synopsis of biological data on the Blue Crab, *Callinectes sapidus* Rathbun, by Mark R. Millikin and Austin B. Williams. March 1984, 39 p.

2. Development of hexagrammids (Pisces: Scorpaeniformes) in the Northeastern Pacific Ocean, by Arthur W. Kendall, Jr., and Beverly Vinter. March 1984, 44 p.

3. Configurations and relative efficiencies of shrimp trawls employed in southeastern United States waters, by John W. Watson, Jr., Ian K. Workman, Charles W. Taylor, and Anthony F. Serra. March 1984, 12 p.

4. Management of northern fur seals on the Pribilof Islands, Alaska, 1786-1981, by Alton Y. Roppel. April 1984, 26 p.

5. Net phytoplankton and zooplankton in the New York Bight, January 1976 to February 1978, with comments on the effects of wind, Gulf Stream eddies, and slope water intrusions, by Daniel E. Smith and Jack W. Jossi. May 1984, 41 p.

6. Ichthyoplankton survey of the estuarine and inshore waters of the Florida Everglades, May 1971 to February 1972, by L. Alan Collins, and John H. Finucane. July 1984, 75 p.

7. The feeding ecology of some zooplankters that are important prey items of larval fish, by Jefferson T. Turner. July 1984, 28 p.

8. Proceedings of the International Workshop on Age Determination of Oceanic Pelagic Fishes: Tunas, Billfishes, and Sharks, by Eric D. Prince (convener and editor), and Lynn M. Pulos (editor). December 1983, 211 p.

9. Sampling statistics in the Atlantic menhaden fishery, by Alexander J. Chester. August 1984, 16 p.

10. Proceedings of the Seventh U.S.-Japan Meeting on Aquaculture, Marine Finfish Culture, Tokyo, Japan, October 3-4, 1978, by Carl J. Sindermann (editor). August 1984, 31 p.

11. Taxonomy of North American fish Eimeriidae, by Steve J. Upton, David W. Reduker, William L. Current, and Donald W. Duszynski. August 1984, 18 p.

12. Soviet-American Cooperative Research on Marine Mammals. Volume 1—Pinnipeds, by Francis H. Fay, and Gennadii A. Fedoseev (editors). September 1984, 104 p.

13. Guidelines for reducing porpoise mortality in tuna purse seining, by James M. Coe, David B. Holts, and Richard W. Butler. September 1984, 16 p.

14. Synopsis of biological data on shortnose sturgeon, *Acipenser brevirostrum* LeSueur 1818, by Michael J. Dadswell, Bruce D. Taubert, Thomas S. Squiers, Donald Marchette, and Jack Buckley. October 1984, 45 p.

15. Chaetognatha of the Caribbean sea and adjacent areas, by Harding B. Michel. October 1984, 33 p.

16. Proceedings of the Ninth and Tenth U.S.-Japan Meetings on Aquaculture, by Carl J. Sindermann (editor). November 1984, 92 p.

17. Identification and estimation of size from the beaks of 18 species of cephalopods from the Pacific Ocean, by Gary A. Wolff. November 1984, 50 p.

18. A temporal and spatial study of invertebrate communities associated with hard-bottom habitats in the South Atlantic Bight, by E. L. Wenner, P. Hinde, D. M. Knott, and R. F. Van Dolah. November 1984, 104 p.

19. Synopsis of biological data on spottail pinfish, *Diplodus holbrooki* (Pisces: Sparidae), by George H. Darcy. January 1985, 11 p.

20. Ichthyoplankton of the Continental Shelf near Kodiak Island, Alaska, by Arthur W. Kendall, Jr., and Jean R. Dunn. January 1985, 89 p.

21. Annotated bibliography on hypoxia and its effects on marine life, with emphasis on the Gulf of Mexico, by Maurice L. Renaud. February 1985, 9 p.

22. Congrid eels of the eastern Pacific and key to their Leptocephali, by Solomon N. Raju. February 1985, 19 p.

23. Synopsis of biological data on the pinfish, *Lagodon rhomboides* (Pisces:Sparidae), by George H. Darcy. February 1985, 32 p.

24. Temperature conditions in the cold pool 1977-81: A comparison between southern New England and New York transects, by Steven K. Cook. February 1985, 22 p.

25. Parasitology and pathology of marine organisms of the world ocean, by William J. Hargis, Jr. (editor). March 1985, 135 p.

26. Synopsis of biological data on the sand perch, *Diplectrum formosum* (Pisces: Serranidae), by George H. Darcy. March 1985, 21 p.

27. Proceedings of the Eleventh U.S.-Japan Meeting on Aquaculture, Salmon Enhancement, Tokyo, Japan, October 19-20, 1982, by Carl J. Sindermann (editor). March 1985, 102 p.

28. Review of geographical stocks of tropical dolphins (*Stenella* spp. and *Delphinus delphis*) in the eastern Pacific, by William F. Perrin, Michael D. Scott, G. Jay Walker, and Virginia L. Cass. March 1985, 28 p.

29. Prevalence, intensity, longevity, and persistence of *Anisakis* sp. larvae and *Lacistorhynchus tenuis* metacestodes in San Francisco striped bass, by Mike Moser, Judy A. Sakanari, Carol A. Reilly, and Jeannette Whipple. April 1985, 4 p.

30. Synopsis of biological data on the pink shrimp, *Pandalus borealis* Kröyer, 1838, by Sandra E. Shumway, Herbert C. Perkins, Daniel F. Schick, and Alden P. Stickney. May 1985, 57 p.

31. Shark catches from selected fisheries off the U.S. east coast, by Emory D. Anderson, John G. Casey, John J. Hoey, and W. N. Witzell. July 1985, 22 p.

32. Nutrient Distributions for Georges Bank and adjacent waters in 1979, by A. F. J. Draxler, A. Matte, R. Waldhauer, and J. E. O'Reilly. July 1985, 34 p.

33. Marine flora and fauna of the Northeastern United States. Echinodermata: Echinoidea, by D. Keith Serafy and F. Julian Fell. September 1985, 27 p.

34. Additions to a revision of the shark genus *Carcharhinus*: Synonymy of *Aprionodon* and *Hypoprion*, and description of a new species of *Carcharhinus* (Carcharhinidae), by J. A. F. Garrick. November 1985, 26 p.

35. Synoptic review of the literature on the Southern oyster drill *Thais haemastoma floridana*, by Philip A. Butler. November 1985, 9 p.

NMFS Honolulu Library
00000000004734

NOAA Technical Report NMFS 44
FAO Fisheries Synopsis No. 148

Synopsis of Biological Data on the Porgies, *Calamus arctifrons* and *C. proridens* (Pisces: Sparidae)

George H. Darcy

September 1986

U.S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
National Oceanic and Atmospheric Administration
Anthony J. Calio, Administrator
National Marine Fisheries Service
William G. Gordon, Assistant Administrator for Fisheries

Library
NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Introduction 1

1 Identity
- 1.1 Nomenclature 1
  - 1.11 Valid name 1
  - 1.12 Objective synonymy 1
- 1.2 Taxonomy
  - 1.21 Affinities 1
  - 1.22 Taxonomic status 3
  - 1.23 Subspecies 3
  - 1.24 Standard common names, vernacular names 3
- 1.3 Morphology
  - 1.31 External morphology 3
  - *1.32 Cytomorphology
  - *1.33 Protein specificity

2 Distribution
- 2.1 Total area 4
- 2.2 Differential distribution 5
  - 2.21 Spawn, larvae, and juveniles 5
  - 2.22 Adults 5
- 2.3 Determinants of distribution changes 5
- 2.4 Hybridization 7

3 Bionomics and life history
- 3.1 Reproduction 7
  - 3.11 Sexuality 7
  - 3.12 Maturity 8
  - 3.13 Mating 8
  - 3.14 Fertilization 8
  - 3.15 Gonads 8
  - 3.16 Spawning 8
  - 3.17 Spawn 9
- 3.2 Preadult phase 9
  - 3.21 Embryonic phase 9
  - 3.22 Larvae and adolescent phase 9
- 3.3 Adult phase 9
  - 3.31 Longevity 9
  - 3.32 Hardiness 9
  - 3.33 Competitors 9
  - 3.34 Predators 9
  - 3.35 Parasites, diseases, injuries, and abnormalities 9
  - *3.36 Chemical composition
- 3.4 Nutrition and growth 9
  - 3.41 Feeding 9
  - 3.42 Food 10
  - 3.43 Growth rate 10
  - 3.44 Metabolism 10
- 3.5 Behavior 10
  - 3.51 Migrations and local movements 10
  - 3.52 Schooling 11
  - *3.53 Responses to stimuli

4 Population
- 4.1 Structure 11
  - 4.11 Sex ratio 11
  - *4.12 Age composition
  - 4.13 Size composition 11
  - 4.14 Subpopulations 14
- 4.2 Abundance and density 14
- 4.3 Natality and recruitment 16
  - 4.31 Reproduction rates 16
  - 4.32 Factors affecting reproduction 16
  - 4.33 Recruitment 16
- *4.4 Mortality and morbidity 16
- *4.5 Dynamics of population 16
- 4.6 The population in the community and the ecosystem 16

5 Exploitation
- 5.1 Fishing equipment 16
- 5.2 Fishing areas 16
- 5.3 Fishing seasons 17
- 5.4 Fishing operations and results 17
  - *5.41 Effort and intensity
  - *5.42 Selectivity
  - 5.43 Catches 17

6 Protection and management
- 6.1 Regulatory measures 17

*7 Pond fish culture

Acknowledgments 17

Citations 18

*No information available.

Synopsis of Biological Data on the Porgies, *Calamus arctifrons* and *C. proridens* (Pisces: Sparidae)

GEORGE H. DARCY
Southeast Fisheries Center
National Marine Fisheries Service, NOAA
75 Virginia Beach Drive
Miami, FL 33149-1099

ABSTRACT

Information on the biology and fishery resources of two common species of western Atlantic porgies, *Calamus arctifrons* and *C. proridens*, is compiled, reviewed, and analyzed in the FAO species synopsis style.

INTRODUCTION

Porgies are small to moderately large spiny-rayed fishes of the family Sparidae. They inhabit temperate and tropical marine and estuarine waters from shore to depths of over 100 m. They are ecologically similar to snappers (Lutjanidae) and grunts (Haemulidae). Most are excellent food-fishes and are of considerable commercial importance (Randall and Vergara R. 1978). Many are also important to recreational fishermen.

The genus *Calamus* consists of about 13 species, all of tropical and subtropical American waters. Two species occur in the eastern Pacific, and 11 occur in the western Atlantic (Randall and Caldwell 1966). Eight species are known from coastal waters of the United States (Robins et al. 1980).

The grass porgy, *Calamus arctifrons*, and the littlehead porgy, *C. proridens*, are two common porgies of the western Atlantic Ocean. The grass porgy is essentially restricted to the northeastern Gulf of Mexico and is fairly abundant in shallow water off the northwest coast of Florida. The littlehead porgy occurs on the east coast of Florida, in the northeastern Gulf of Mexico, in the Greater Antilles, and on Campeche Bank where it is one of the most important commercial fishes. This paper summarizes information on these species.

1 IDENTITY

1.1 Nomenclature

1.11 Valid name

Grass porgy, *Calamus arctifrons* Goode and Bean, 1882 (Fig. 1): 425, type locality: Pensacola, Florida. The name comes from the Latin *calamus* (a reed or quill), referring to the quill-like interhaemal bone (Randall and Caldwell 1966), and *arctus* (narrow) and *frons* (forehead), referring to the dorsal profile of the head.

Littlehead porgy, *Calamus proridens* Jordan and Gilbert, 1884 (Fig. 2): 150, type locality: Key West, Florida. The name comes from the Latin *calamus* (see above) and *prora* (prow) and *dens* (tooth), referring to the projecting canine teeth (Jordan and Fesler 1893).

1.12 Objective synonymy

Calamus arctifrons Goode and Bean, 1882

Calamus proridens Jordan and Gilbert, 1884

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Class Osteichthyes
Superorder Acanthopterygii
Order Perciformes
Suborder Percoidei
Family Sparidae



Figure 1.—Adult *Calamus arctifrons*. (From Jordan and Evermann 1896, fig. 550)



Figure 2.—Adult *Calamus proridens*. (From Jordan and Evermann 1896, fig. 547)

Generic

Systematics of the genus *Calamus* have been badly confused in the past (Randall and Caldwell 1966). Reviews of the genus have appeared in Valenciennes in Cuvier and Valenciennes (1830), Guichenot in de la Sagra (1843), Guichenot (1868), Poey (1874), Jordan and Gilbert (1882), Jordon and Fesler (1893), Jordan and Evermann (1898), Evermann and Marsh (1902), Longley and Hildebrand (1941), and Randall and Caldwell (1966). Species are often difficult to differentiate, and are especially difficult to identify when small. According to Randall and Caldwell (1966), 13 species of *Calamus* exist. All are American, with two species from the eastern Pacific, and 11 from the western Atlantic.

The genus is characterized as follows: Second interhaemal bone enlarged and hollowed anteriorly, receiving posterior portion of swimbladder; dorsal fin preceded by a small antrorse (forward-projecting) spine, often hidden by scales; temporal crests thin and high, joining lateral crests, which form part of the orbit above middle of eye, both crests coalescing with supraoccipital in narrow anterior part of interorbital; body deep, the depth contained 1.8-2.5 times in standard length (SL), and compressed, the width about 6 in SL; head large, its length 2.8-3.6 in SL; dorsal profile of head from moderately to very steep; eye elevated, the suborbital depth 5.4-12.5 in SL (often greater in adults than in juveniles); mouth terminal, not large; anterior teeth in jaws conical, those on sides molariform, in two full rows in lower jaw and three rows in upper jaw (teeth in middle row small); a bony tubercle projecting downward from lower anterior portion of maxillaries; prefrontal with a bony tubercle above posterior nostril, this tubercle much more conspicuous in some species than in others; posterior nostril long and slit-like; preopercle and opercle entire, the margins smooth; dorsal fin rays XII (rarely XI or XIII), 11 or 12 (rarely 13); anal fin rays III, 10 or 11 (rarely 9); pectoral fin rays 14-16 (rarely 13 or 17); pelvic fin rays I, 5; 15 branched caudal fin rays; body completely covered with weakly-ctenoid scales; head with scales on opercle, preopercle, interopercle, and extending anteriorly from nape into interorbital region; no scales on dorsal and anal fins except basally on last two rays; scales extend about two-thirds of way onto caudal fin and occur basally on paired fins, dorsal fin slightly notched between spinous and soft portions; dorsal fin relatively low, the longest spine (third or fourth) contained 5.6-9.6 in SL; pectoral fin length 2.1-3.6 in SL; caudal fin forked; gill rakers short, 9-14 on first gill arch; vertebrae 10+14; three predorsal bones; dorsal fin support count 0-0-0-2 (Randall and Caldwell 1966). The skull was discussed by Jordan and Gilbert (1882), and illustrated by Gregory (1933).

Specific

The most important characters for separating species of *Calamus* are: Number of lateral-line scales; total number of pectoral fin rays; number of dorsal and anal fin rays; dentition; eye diameter; suborbital depth; length of pectoral fins; length of the longest dorsal spine (usually the third spine); body depth; color; and steepness of profile of the head (Randall and Caldwell 1966).

Calamus arctifrons—The following species diagnosis is from Randall and Caldwell (1966): Dorsal fin rays XI to XIII (usually XII), 11 to 13 (usually 12); anal fin rays III, 10 or 11 (usually 10); pectoral fin rays usually 16 (sometimes 15); lateral-line scales 43-49 to base of caudal fin; total gill rakers in first gill arch 10-12 (usually 10); canine teeth in front of jaws about equal in size; no small molariform teeth medial to the three rows at sides of upper jaw; pectoral fins short, their length 3.0-3.6 in SL; eye small; suborbital depth moderately great, 6.6-7.8 in SL; body depth 2.0-2.5 in SL (depth relatively less in adults than juveniles); longest dorsal spine 6.4-8.9 in SL; dorsal profile of head usually smoothly convex, the portion below eye forming an angle of about 50-57° to the horizontal (taken from tip of snout above upper lip to mid-base of caudal fin); maxillary tubercle with a free semicircular margin; prefrontal tubercle not well developed.

Calamus arctifrons is very similar to *C. campechanus*, and probably evolved from a common stock in the not too distant past (Randall and Caldwell 1966). It is distinguished from *C. campechanus* by its smaller suborbital depth, higher number of pectoral fin rays, lower number of gill rakers, and lower dorsal profile of the head below level of the eye (Randall and Caldwell 1966). All specimens of *C. campechanus* examined by Randall and Caldwell had irregular horizontal lines on the unscaled portion of the cheeks, whereas only a few specimens of *C. arctifrons* had these lines. The two species are separated geographically, with *C. campechanus* restricted to Campeche Bank. Environmental conditions such as tubidity, temperature, and salinity apparently exclude both species from the western Gulf of Mexico (Randall and Caldwell 1966).

Calamus proridens—The following species diagnosis is from Randall and Caldwell (1966): Dorsal fin rays XII or XIII (usually XII), 11 or 12 (usually 12); anal fin rays III, 10 or 11 (usually 10); pectoral fin rays 13-15 (usually 14); lateral-line scales 52-57 (usually 54-56); total gill rakers in first gill arch 11-13 (usually 11); fourth canine tooth from symphysis of upper jaw enlarged and strongly outcurved in adults (enlargement begins at about 100 mm SL and is apparent at sizes at least as small as 148 mm SL); a group of small, molariform teeth medial and toward the front of the three rows at sides of upper jaw, forming a partial fourth row at front half of sides of jaw; pectoral fins long, 2.5-3.0 in SL; eye small; suborbital depth relatively great; body depth 1.95-2.20 in SL; longest dorsal fin spine 6.1-8.7 in SL (in specimens >100 mm SL); prefrontal tubercle above posterior nostril not well developed; largest molariform tooth usually smaller than fleshy width of maxillary tubercle; dorsal profile of head steep and relatively straight, portion below eye of adults forming an angle of 58-65° from horizontal and first third above eye an angle of 43-69°.

Calamus proridens is closely related to *C. pennatula*, but differs in having a steeper profile of the upper part of the head, greater average body depth, greater average length of the longest dorsal fin spine (third or fourth) and slightly higher lateral-line scale counts (Randall and Caldwell 1966). Large specimens of *C. proridens* develop a large, bulging forehead not found in *C. pennatula*.

1.22 Taxonomic status

Both *C. arctifrons* and *C. proridens* are considered morphospecies. *Calamus arctifrons* is very closely related to *C. campechanus*, but they are considered separate species (Randall and Caldwell 1966; Randall and Vergara R. 1978).

See 1.21.

1.23 Subspecies

No subspecies of *C. arctifrons* or *C. proridens* are recognized.

1.24 Standard common names, vernacular names

Calamus arctifrons—The accepted common name for *C. arctifrons* in the United States is grass porgy (Robins et al. 1980), and standard FAO common names are: English, grass porgy; French, daubenet cendre; Spanish, pluma negra (Randall and Vergara R. 1978). Other names appearing in the literature are shad porgy (Jordan and Fesler 1893; Evermann and Marsh 1902; Beebe and Tee-Van 1928), pez de pluma (Duarte-Bello 1959) and porgy (Springer and Woodburn 1960). According to Moe (1963), *C. arctifrons*, *C. proridens*, and *C. calamus* are caught on the snapper grounds off the west coast of Florida, and the names bank porgy, Key West porgy, grass porgy, and white snapper are applied to all three species, although Key West porgy usually refers to *C. nodosus*.

Calamus proridens—The accepted common name of *C. proridens* in the United States is littlehead porgy (Robins et al. 1980), and standard FAO common names are: English, littlehead porgy; French, daubenet titête; Spanish, pluma joroba (Randall and Vergara R. 1978). Other names or variations appearing in the literature include little-head porgy (Jordan and Fesler 1893; Beebe and Tee-Van 1928), pez de pluma (Jordan and Fesler 1893; Beebe and Tee-Van 1928; Duarte-Bello 1959), bajonado (Campeche Bank) (Sokolova 1965), crocro and tête-feuilles (Trinidad) (Zaneveld 1983) and bajonado azul (Campeche Bank) (Juárez 1975). The names bank porgy, Key West porgy, grass porgy, and white snapper are sometimes applied indiscriminately to *C. proridens*, *C. arctifrons*, and *C. calamus* from the snapper grounds off the west coast of Florida (Moe 1963).

1.3 Morphology

1.31 External morphology

Calamus arctifrons—The following morphological description is adapted from Evermann and Marsh (1902) and Randall and Vergara R. (1978): Head length 3.3 in SL; body oval, compressed and deep, its depth 2.0-2.5 in SL; eye width 3.4 in SL; snout length 1.9 in SL; maxillary length 2.4 in SL; interorbital distance 3.5 in SL; preorbital distance 2.8 in SL; suborbital space rather deep, 6.6-7.8 in SL; upper head profile usually smoothly convex and moderately steep; jaws anteriorly with about 8-10 canine-like teeth of about equal size; jaws with two rows of molar-like teeth laterally in lower jaw and three rows in upper jaw, without an irregular series inside and toward the front; dorsal fin XII, 12; anal fin III, 10; pectoral fins relatively short, not reaching to anal fin origin when appressed, usually with 16 rays; lateral-line scales 43-49; about five rows of scales on cheek; back not greatly elevated; anterior profile not steep, a slight angle in front of eye.

Additional meristic data are presented in Table 1. The skull of *Calamus* sp. was discussed by Jordan and Gilbert (1882), and illustrated by Gregory (1933).

Table 1.—Meristic data for *Calamus arctifrons*. Numbers in parentheses are percentages of the total number of specimens examined. (From Randall and Caldwell 1966)

Character	Count (%)						
Lateral-line scales	43 (1.2)	44 (8.6)	45 (9.9)	46 (32.1)	47 (29.6)	48 (9.9)	49 (8.6)
Pectoral fin rays			15-15 (11.5)	15-16 (20.5)	16-16 (67.9)		
Dorsal fin spines	XI		XII			XIII	
Dorsal fin rays	12 (1.3)	13 (0)	11 (5.2)	12 (85.7)	13 (1.3)	11 (6.5)	12 (0)
Anal fin spines			III				
Anal fin rays			9 (0)	10 (97.4)	11 (2.6)		
Gill rakers in first arch			10 (63.8)	11 (27.5)	12 (8.7)		

Color of adults in preservative: Light brown; a black spot smaller than the eye, but larger than the pupil near the front of the lateral line; five vertical rows of dark brown blotches on body and two diagonally elongate blotches on caudal peduncle, the last continuing as the upper part of a large V-shaped marking basally on caudal fin; two diffuse dark blotches mid-dorsally on nape anterior to first row of spots on body; occasional specimens have irregular horizontal lines on the unscaled portions of cheeks; dorsal fins with some faint dark blotches in line with rows of blotches on side of body; lobes of caudal fin traversed by about five blackish bands; pelvic fins with two faint transverse dark blotches (Randall and Caldwell 1966).

Color of juveniles in preservative: Darker rows of blotches than in adults, and a dark bar on head running through the eye, its front

edge touching opening of the mouth; a few dark streaks on opercle and scattered brown dots on body among the dark blotches forming the dark bars; two dark spots dorsally on caudal peduncle; a row of dark spots on dorsal fin with a second row in spinous portion above and a second row in soft portion below; two rows of dark spots on anal fin; dark bars on caudal lobes very conspicuous; pelvic fins with two broad, dark crossbars (Randall and Caldwell 1966).

Description of colors in life: Silvery, olivaceous or bluish above, the centers of many of the scales pearly and iridescent; paler below; back and sides with 6-8 dark vertical crossbars, narrower than interspaces, with spots between them; conspicuous black blotch on lateral line near upper end of gill opening; row of about six salmon-olive spots along lateral line; snout olive, mottled with bluish and may be streaked with yellow; interorbital area may have a yellow band; broad, dark bar from eye straight downward across cheek; an obscure pearly-blue streak below eye, and two or three similar ones before eye; preorbital usually bluish, with more or less numerous longitudinal streaks and dashes of golden yellow; preorbital sometimes pale salmon-yellow, with a few light bluish streaks; cheeks, preopercle, and opercle pearly, with yellow shades and spots; opercular membrane coppery orange; pectoral fins pale; pelvic fins bluish white, faintly barred; dorsal fin bluish, with blackish spots along base of dorsal; anal fin bluish; caudal fin bluish, with small, dusky salmon spots forming undulating crossbars, posterior margin slightly dusky (Goode and Bean 1882; Jordan and Gilbert 1884; Evermann and Marsh 1902; Randall and Vergara R. 1978).

See 1.21.

Calamus proridens—The following morphological description is adapted from Randall and Caldwell (1966) and Randall and Vergara R. (1978): Body oblong, compressed, very deep anteriorly (1.95-2.20 in SL, relatively less in adults than in juveniles); snout blunt and very steep in adults (less so in juveniles); dorsal profile of head usually smoothly convex; nape strongly convex, developing into a distinct bump in large specimens; suborbital space rather deep, about 6.3-9.9 in SL; mouth small (especially in small specimens), the maxilla not reaching to anterior eye margin; eye small; both jaws anteriorly with canine-like teeth, the fourth from midline on each side enlarged and noticeably outcurved in specimens >18 cm SL; laterally with molar-like teeth in three rows plus an irregular series inside and toward the front in upper jaw; dorsal fin rays XII or XIII (usually XII), 11 or 12 (usually 12); anal fin rays III, 10 or 11 (usually 10); pectoral fin rays 13-15 (usually 14), the pectoral fins extending to or beyond anal fin origin when appressed; lateral-line scales 52-57 (usually 54-56); total gill rakers in first gill arch 11-13 (usually 11).

Additional meristic data are presented in Table 2. The skull of *Calamus* sp. was discussed by Jordan and Gilbert (1882), and illustrated by Gregory (1933).

Color in preservative: Tan to light brown; center of each scale on upper half of body with a dark brown spot; horizontally elongate dark brown blotch, broader posteriorly, running over upper end of gill opening; front end of blotch beginning at or slightly anterior to dorsal end of free margin of preopercle; supra and suborbital dark brown lines; three dark brown lines extending anteriorly from region in front of eye between supra and suborbital lines; unscaled suborbital region with well-defined dark brown horizontal or slightly diagonal lines on a bronze background, usually slightly wavy and sometimes broken into spots or lines; caudal fin about same color as body; other fins paler, dorsal with some faint brownish blotches (Randall and Caldwell 1966).

Table 2.—Meristic data for *Calamus proridens*. Numbers in parentheses are percentages of the total number of specimens examined. (From Randall and Caldwell 1966)

Character	Count (%)					
Lateral-line scales	52 (2.7)	53 (5.4)	54 (29.7)	55 (37.8)	56 (18.9)	57 (5.4)
Pectoral fin rays		13-14 (2.7)	13-15 (0)	14-14 (83.8)	14-15 (13.5)	
Dorsal fin spines		XII		XIII		
Dorsal fin rays		11 (2.8)	12 (94.4)	11 (2.8)	12 (0)	
Anal fin spines		III				
Anal fin rays			10 (91.9)	11 (8.1)		
Gill rakers in first arch		11 (75.9)	12 (20.7)	13 (3.4)		

Color in life: Iridescent silvery, with bright bluish tinges on back and upper sides; diffuse horizontal elongate blue blotch at upper end of gill opening; blue streak running along lower eye margin and alternating blue (narrow) and yellow (wide) horizontal lines across unscaled portion of cheeks; unmarked portion of suborbital and snout brassy; lips yellowish; inside corners of mouth yellow (Randall and Caldwell 1966; Randall and Vergara R. 1978).

See 1.21.

2 DISTRIBUTION

2.1 Total area

Calamus arctifrons—*Calamus arctifrons* occurs in the northeastern Gulf of Mexico, primarily in shallow water along the west coast of Florida, but straying on occasion to Louisiana and Texas (Fig. 3) (Randall and Caldwell 1966; Hoese and Moore 1977; Randall and Vergara R. 1978). The species is apparently absent from the northwestern Gulf of Mexico (Briggs 1958; Randall and Caldwell 1966), and Campeche Bank. Although there are several published reports of *C. arctifrons* from the West Indies (Evermann and Marsh 1902, Puerto Rico; Beebe and Tee-Van 1928, Haiti), Randall and Caldwell (1966) were not able to find any West Indian specimens in museum collections; if the species does occur in the Greater Antilles, it is apparently quite uncommon there. Briggs (1958) placed the southern limit of the range at Ilha Grande, Brazil, which is almost certainly incorrect.

Calamus proridens—*Calamus proridens* occurs along the Atlantic and Gulf of Mexico coasts of Florida to about Louisiana, on Campeche Bank, and through the Greater Antilles (Fig. 4) (Briggs 1958; Randall and Caldwell 1966; Dubovitsky 1977a; Randall and Vergara R. 1978). This species, like *C. arctifrons*, is apparently absent from the northwestern Gulf of Mexico (Briggs 1958; Walls 1975). Struhsaker (1969) reported *C. proridens* as rare, but present, on live bottom in the South Atlantic Bight. Although several authors (Jordan and Gilbert 1884; Jordan and Fesler 1893; Beebe and Tee-Van 1928) reported specimens from the West Indies, Randall and Caldwell (1966) speculated that at least some of these reports may not have been based on natural occurrences, the fish having been caught at some other location. *Calamus proridens* is fished in Cuba (Randall and Vergara R. 1978), and is known to occur in Hispaniola.



Figure 3.—Distribution of *Calamus arctifrons*. (Based on Randall and Vergara R. 1978)



Figure 4.—Distribution of *Calamus proridens*. (Based on Randall and Vergara R. 1978)

2.2 Differential distribution

2.21 Spawn, larvae and juveniles

Calamus arctifrons—Spawning areas of *C. arctifrons* are unknown. Unidentified sparid larvae collected by Houde et al. (1979) in the northeastern Gulf of Mexico probably included *C. arctifrons*. Most (>90%) of the unidentified sparid larvae were collected at stations <50 m deep, and were most common in winter and spring.

Juveniles are often found in shallow seagrass beds. Most of the specimens examined by Tabb and Manning (1961) from Florida Bay *Thalassia* beds were juveniles of 7-10 cm SL.

See 2.3 and 3.16.

Calamus proridens—Spawning areas of *C. proridens* along the coast of the United States are not well known. Unidentified sparid larvae collected by Houde et al. (1979) in the northeastern Gulf of Mexico probably included *C. proridens* (see above).

On Campeche Bank, Dubovitsky (1977a) reported a steady drift of pelagic eggs and larvae westward across the bank, eventually reaching shallow water where the demersal habit is adopted.

Juveniles are found in relatively shallow water. On Campeche Bank, young stay in shallow, productive waters for the first year of life (Dubovitsky 1977a). Between ages I and II, there is a gradual movement offshore toward the northeast and east, where small fish become recruited to the commercial stock.

See 2.3, 3.16, and 3.51.

2.22 Adults

Calamus arctifrons—Adult *C. arctifrons* occur on shallow seagrass beds and other vegetated bottom (Evermann and Marsh 1902; Tabb and Manning 1961; Vick 1964; Randall and Caldwell 1966; Walls 1975; Hoese and Moore 1977; Randall and Vergara R. 1978), and over hard or rock bottoms (Randall and Caldwell 1966; Darcy and Gutherz[3]). They occur from near shore to at least 29.6 m (Darcy and Gutherz[3]), and are demersal.

See 2.1, 2.3, and 3.51.

Calamus proridens—Adult *C. proridens* occur in shallow water, from nearshore to at least 59.3 m on the West Florida Shelf (Darcy and Gutherz unpubl.), and at least 85 m on Campeche Bank (Dubovitsky 1977a). They are found on or near natural and artificial reefs (Smith et al. 1979; Miller and Richards 1980), around offshore platforms (Hastings et al. 1976), and on live bottoms of sponges and corals (Struhsaker 1969; Darcy and Gutherz unpubl.). Smith (1976) termed *C. proridens* a secondary reef fish.

See 2.1, 2.3, and 3.51.

2.3 Determinants of distribution changes

Calamus arctifrons—Distribution of *C. arctifrons* is affected by temperature, salinity, water clarity, water depth, and habitat. In the northern part of its range, *C. arctifrons* appears to be uncommon or absent in shallow water in the coldest months. At Crystal River, Florida, Reid (1954) collected *C. arctifrons* from June through December (except for August); Grimes (1971) collected specimens March through December 1969, with greatest abundance in September and July; and Grimes and Mountain (1971) collected specimens April through December 1970, with greatest abundance in October and December. In the southern part of its range, *C. arctifrons* may remain in shallow water in winter; Tabb and Manning (1961) found them most common October to April in Florida Bay. The species occurs in water of at least 7.8-32.2°C (Crystal River, Reid 1954), and at salinities of at least 9.7-35.7‰ (Crystal River, Reid 1954; Charlotte Harbor, Wang and Raney 1971). *Calamus arctifrons* prefers clear water (Tabb and Manning 1961). Turbidity avoidance may be a reason why this species is absent from the northwestern Gulf of Mexico (Randall and Caldwell 1966).

Calamus arctifrons is stricted to fairly shallow water, perhaps because benthic vegetation is confined to shallow water, and *C. arctifrons* is often associated with vegetation (Evermann and Marsh 1902; Tabb and Manning 1961; Walls 1975; Randall and Vergara R. 1978). The species has been reported from at least as deep as 29.6 m (Darcy and Gutherz unpubl.), but is usually most common in shallower water. At Crystal River, Reid (1954) found *C. arctifrons* on shallow and deep flats and in channels 4.8-5.5 deep. On

the West Florida Shelf, Darcy and Gutherz (unpubl.) reported *C. arctifrons* most common in 9-18 m of water, with a mean depth of 15.8 m, in the surveyed area. In addition to vegetated bottom, *C. arctifrons* is common on hard bottom (Randall and Caldwell 1966) and rock bottom (Darcy and Gutherz unpubl.).

See 2.1, 2.21, 2.22, and 3.51.

Calamus proridens—Distribution of *C. proridens* is affected by salinity, temperature, water depth, growth stage of the fish, and habitat. Although there are no specific references in the literature to salinities at which *C. proridens* occurs, it is not found in estuaries or low salinity area, and probably requires a salinity of at least 30‰.

Distributional changes with depth and season are probably attributable to temperature changes. Little is known about the distribution of *C. proridens* on the Atlantic coast of the United States. In the Gulf of Mexico, the species occurs from at least 11 to 59 m (Darcy and Gutherz unpubl.), with greatest abundance in 19-27 m (in January) and a mean depth occurrence of 27.8 m. Abundance was found to vary significantly with depth. Smith (1976) reported *C. proridens* from 29-42 m depths on the Florida Middle Ground.

On Campeche Bank, considerable study has been made of seasonal distribution changes of *C. proridens*. Although Olaechea and Sauskan (1974) did not feel that water temperature influenced abundance of *Calamus* spp. on Campeche Bank, most other authors (Gonzáles 1973; Olaechea 1975; Dubovitsky 1977a) did report relationships between bottom temperature and *C. proridens* abundance. Gonzáles (1973) found good correlation between catch rates of *Calamus* spp. and water temperature on the eastern edge of the bank, with fish moving deeper on the slope when shallow waters warmed. Olaechea and Sauskan (1974) estimated highest abundance of *C. proridens* on the bank in spring and summer. Sauskan and Olaechea (1974) reported this species most common in shallow (15-30 m) water on the central part of the bank, but also common in shallow and deep (50-120 m) water on the eastern bank, and at moderate (30-50 m) depths on the western and central bank (see section 4.2).

Dubovitsky (1977a) presented detailed information on seasonal changes in geographical distribution of *C. proridens* on Campeche Bank. In January (Fig. 5A), the bulk of the population was on the central bank. When bottom temperatures on the bank did not exceed 22°C, *C. proridens* extended its distribution into deep (65-70 m) water and as far south as lat. 20°N. In February and March (Fig. 5B), *C. proridens* was most common in coastal areas of the eastern bank at depths to 25 m, but extended to 40 m where cold water flowed onto the edge of the shelf. In May through June (Fig. 6), *C. proridens* was restricted to shallow water, with commercially exploitable aggregations present. By August (Fig. 7A), fish spread out and moved to somewhat deeper (60-85 m) water; they were most abundant east of long. 89°30′W, and below 30 m depth. Bottom waters from May to August are the coldest of the year on the bank, due to upwelling. In November, bottom temperatures are at their maximum for the year, and fish were found in fairly shallow water (Fig. 7B), with large aggregations only in the eastern and western parts of the bank. In December (Fig. 7B), a similar pattern was seen, with fish well spread out across the bank. In general, aggregations formed in periods of lowest bottom temperatures.

Seasonal changes in size composition of *C. proridens* also occur on Campeche Bank (Dubovitsky 1977a). From December to March, large individuals were always present on the eastern bank where bottom waters were cold, though the numbers of large individuals decreased from December to March. Large fish were present in shallow (<30-35 m) water on the central bank in December, small fish were present in January, and both size groups were present in March. On the offshore part of the bank, at 35-55 m, small (18-20 cm, type of length measurement unspecified) fish dominated from December to March, their numbers increasing with depth. Juveniles of 13-16 cm appeared in offshore waters of the western and central bank in January, and on the eastern bank in February. From May through August, size distributions were similar to those of January. In November, small fish dominated everywhere except on the shallow central bank, where mixed size groups occurred, and on the northwestern bank. There was a notable increase in small, immature fish on the southwestern bank. In general, throughout the year, large fish were associated with low bottom temperatures and were most common on the eastern bank where cold-water upwelling occurred. Large fish penetrated the western bank only when bottom temperatures there declined. Small fish were usually associated with higher bottom temperatures and were most common on the western bank and in shallow water, extending eastward only in times of high bottom temperatures.



Figure 5.—Distribution of *Calamus proridens* and catch rates on Campeche Bank; A. January 1970; B. February-March 1971. (From Dubovitsky 1977a, fig. 2)



Figure 6.—Distribution of *Calamus proridens* and catch rates on Campeche Bank; A. May 1969; B. June 1969. (From Dubovitsky 1977a, fig. 2)



Figure 7.—Distribution of *Calamus proridens* and catch rates on Campeche Bank; A. August 1969 (circles) and August 1971 (triangles); B. November 1970 (circles) and December 1970 (triangles). (From Dubovitsky 1977a, fig. 2)

Calamus proridens is found primarily on sponge and coral bottoms (live bottom) (Struhsaker 1969; Darcy and Gutherz unpubl.). Smith et al. (1979) reported *C. proridens* from natural and artificial reefs off Clearwater, Florida, and Smith (1976) listed it as a secondary reef fish of the northeastern Gulf of Mexico. The species also inhabits offshore platforms in the northern Gulf (Hastings et al. 1976). Hard substrates and sessile invertebrate growths probably provide shelter and harbor food organisms of *C. proridens*.

See 2.1, 2.21, 2.22, 3.51, 3.52, 4.13, and 4.2

2.4 Hybridization

No hybrids of either *C. arctifrons* or *C. proridens* are known.

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

Nothing is known of sexuality of *C. arctifrons*, though it is probably a hermaphroditic species like many other sparids. *Calamus proridens* is known to be a protogynous hermaphrodite, with individuals changing from females to males at about 24 cm (measurement unspecified) (Dubovitsky 1977b). Because of this sex reversal, females are consistently smaller than males (Fig. 8).

See 3.12 and 3.15.



Figure 8.—Mean lengths of males (squares) and females (circles) of *Calamus proridens* from Campeche Bank. Type of length measurement was unspecified. (From Dubovitsky 1977a, fig. 5)

3.12 Maturity

Maturation size and age of *C. arctifrons* are unknown. Although Dubovitsky (1977a, b) did not state a size of maturation for *C. proridens*, he did identify females as small as about 17 cm (measurement unspecified), and males as small as about 23 cm. The great majority of adult females collected by Dubovitsky (1977a) were ripe and in prespawning condition, but never more than half of the ovaries had eggs in ripening condition. Male maturation stages were difficult to assess.

See 3.11 and 3.16.

3.13 Mating

Mating in *C. arctifrons* and *C. proridens* has not been described. Dubovitsky (1977a) stated that *C. proridens* apparently does not form dense spawning aggregations.

3.14 Fertilization

Fertilization is probably external.

3.15 Gonads

Gonads of *C. arctifrons* have not been described. Dubovitsky (1977a, b) histologically examined gonads of *C. proridens*, and found successive sex inversion from female to male. The size at which the ovaries regress and testes develop is very variable, but there is a pronounced sex inversion at about 23-24 cm (measurement unspecified). By the end of the spawning season (middle to late summer), asymmetry of the ovaries is often observed, the left ovary being larger.

See 3.12 and 3.16.

3.16 Spawning

Calamus arctifrons—Spawning season of *C. arctifrons* is not known with certainty. However, unidentified sparid larvae collected in the northeastern Gulf of Mexico by Houde et al. (1979), which probably included *C. arctifrons*, were most abundant in winter and spring. Reid (1954) hypothesized early spring spawning at Crystal River, based on small individuals caught in June and July.

Calamus proridens—Spawning season of *C. proridens* along the coast of the United States is not known, but larval surveys by Houde et al. (1979) in the eastern Gulf of Mexico indicated highest abundance of sparid larvae, probably including *C. proridens*, in winter and spring.

On Campeche Bank, several studies of *C. proridens* have indicated a long spawning period. Juárez (1975) stated that the spawning season lasts from summer into winter, with maximum spawning intensity in winter. Dubovitsky (1977a) reported spawning from December to August, with a peak from January to March. Based on visual inspection and histological examination of ovaries, individuals are apparently partial spawners (Dubovitsky 1977a), and may spawn more than once a year. No dense spawning aggregations have been reported, but localized areas of prespawning (characterized by numerous large, transparent oocytes occupying, visually, about 20% of the total ovary volume) *C. proridens* have been identified (Fig. 9) (Dubovitsky 1977a). Large, ripe individuals occurred from December to June, with greatest abundance in December and February on the shallow eastern bank, and in January on the western bank at depths >35 m. Small prespawning fish were observed from January through August, with peaks in January, March, and July. Ripe, small fish were observed in offshore areas at depths >40-50 m, and only on the central bank. Sequential spawning of size groups was also reported by Dubovitsky (1977a), with different spawning locations for different size groups. Over the spawning season, a gradual diminution of size of spawners occurred. From late May to the beginning of June, average size was 22 cm (measurement unspecified); in early June it was 20 cm; by late July, 18 cm; and by early August, 17 cm.



Figure 9.—Locations and months of collection of *Calamus proridens* in pre-spawning condition on Campeche Bank; A. December to March; B. May to August. (From Dubovitsky 1977a, fig. 6)



Figure 10.—Seasonal changes in reproductive activity of *Calamus proridens* and related factors on Campeche Bank. A. Bottom temperatures, 1969-1971. B. Total plankton biomass between the 20-m and 200-m isobaths for lat. 91°W (dotted line) and lat. 87°20'W (solid line). C. Percentage of pre-spawning females captured. Plankton productivity figures are from Jrómov (1967). (From Dubovitsky 1977a, fig. 7)

Spawning of *C. proridens* on Campeche Bank is temperature-dependent, beginning with the appearance of cold-water inflows onto the eastern bank in December (Dubovitsky 1977a). Plankton productivity is also temperature-dependent, with maximum productivity in January through August (Jrómov 1967). High plankton productivity at the time of spawning assures that larval stages will have adequate food supply. The relationships between temperature, plankton productivity, and spawning of *C. proridens* on Campeche Bank are shown in Fig. 10.

See 2.21, 2.23, 3.51, and 4.13.

3.17 Spawn

Eggs of *C. arctifrons* are probably pelagic, since most sparids produce pelagic eggs with little or no pigment (Breder 1962). Eggs of *C. proridens* are known to be pelagic (Dubovitsky 1977a).

3.2 Preadult phase

3.21 Embryonic phase

Embryonic development of *C. arctifrons* and *C. proridens* have not been studied. Developing eggs are probably pelagic.

3.22 Larvae and adolescent phase

Larvae of *C. arctifrons* and *C. proridens* have not been described in the literature. Young of *Calamus* spp. are difficult to distinguish. Houde et al. (1979) were unable to separate species of *Calamus* from plankton collections in the eastern Gulf of Mexico. *Calamus proridens* is known to have pelagic larvae (Dubovitsky 1977a); *C. arctifrons* larvae are probably also pelagic.

3.3 Adult phase

3.31 Longevity

Calamus arctifrons reaches a maximum size of about 22 cm (measurement unspecified) and is common to 20 cm (Randall and Vergara R. 1978). *Calamus proridens* reaches 44 cm and is common to 37 cm (Randall and Vergara R. 1978). Life spans of these species are unknown.

3.32 Hardiness

Calamus arctifrons is apparently subject to toxic effects of red tides, because Springer and Woodburn (1960) found several dead individuals on the beach in Pinellas County, Florida, following an outbreak.

See 2.3 and 3.35.

3.33 Competitors

Competitors of *C. arctifrons* and *C. proridens* are not known, but probably include other sparids, haemulids, lutjanids, labrids, and other fishes that feed on benthic invertebrates. Associates of *C. arctifrons* and *C. proridens* on the West Florida Shelf are discussed in section 4.6. Whether these associates are actually competitors is not known.

3.34 Predators

Predators of *C. arctifrons* and *C. proridens* are unknown, but probably include sharks, sphyraenids, and large serranids and lutjanids.

3.35 Parasites, diseases, injuries, and abnormalities

Calamus arctifrons has been reported parasitized by intestinal trematodes, including *Proctoeces lintoni*, *Lepocreadium opsanusi*, *Pseudocreadium anandrum*, and *Pachycreadium crassigulum* (Nahhas and Cable 1964). Red tides apparently prove fatal to some individuals of *C. arctifrons* (Springer and Woodburn 1960).

3.4 Nutrition and growth

3.41 Feeding

Feeding of *C. arctifrons* is unknown, but dentition suggests benthic feeding habits. Sessile and motile invertebrates are probably bitten or picked off hard substrates and seagrass blades, and other prey picked from the surface of softer substrates.

Calamus proridens probably feeds in a manner similar to that described for *C. arctifrons*. On Campeche Bank, feeding intensity varies over the year (Fig. 11), with major feeding concentrated around the spawning and postspawning periods (Dubovitsky 1977a). Primary feeding grounds on Campeche Bank are the inshore parts of the plateau (Dubovitsky 1977a). At the outer edge of the bank, intense feeding, as measured by mean index of stomach fullness, was noted only in the period just prior to spawning.

See 3.42.



Figure 11.—Feeding intensity of small mature (circles) and large mature (triangles) specimens of *Calamus proridens* by month on Campeche Bank. Solid symbols refer to specimens collected in >30-35 m depths, open symbols refer to specimens collected in depths <30-35 m. (From Dubovitsky 1977a, fig. 9)



Figure 12.—Seasonal changes in fat content of mature *Calamus proridens* from Campeche Bank. Circles represent small specimens, triangles represent large specimens. Solid symbols refer to specimens caught in depths >30-35 m, open symbols refer to specimens caught at depths <30-35 m. (From Dubovitsky 1977a, fig. 8)

3.42 Food

Both *C. arctifrons* and *C. proridens* probably feed primarily on invertebrates. Beebe and Tee-Van (1928) reported unidentifiable animal and vegetable materials from stomachs of *C. arctifrons*, including *Thalassia* leaves. Whether the leaves were consumed deliberately or incidentally is not known. *Calamus arctifrons* at Crystal River conatined copepods, amphipods, mysids, shrimp, bivalves, and gastropod mollusks (*Mitrella* sp. and *Bittium* sp.) and polychaetes (Reid 1954). *Calamus proridens* examined by Beebe and Tee-Van (1928) contained small crustaceans. Dentition of *C. proridens* suggests feeding on hard-shelled benthic crabs and bivalved mollusks (Dubovitsky 1977a). On the western part of Campeche Bank, south of lat. 20°N, the main food of young *C. proridens* is pink shrimp, *Penaeus duorarum* (Dubovitsky 1977a).

3.43 Growth rate

Growth rate of *C. arctifrons* is unknown. Growth rate of *C. proridens* is reportedly rapid in the first year of life (Dubovitsky 1977a), though Dubovitsky did not publish growth curves. There is apparently substantial variation in timing of growth and spawning within the population, and scale increment formation is variable, possibly due to differences in spawning and feeding among fish of different size groups (Dubovitsky 1977a). Growth rate differs from female to male phase of the fish; egg development reportedly slows growth of individuals functioning as females (Dubovitsky 1977b), after which the growth rate increases again. Dubovitsky based these conclusions on histological examinations of gonads and analysis of length frequencies, but there is little data presented in his publications on which to critically evaluate them.

See 3.44.

3.44 Metabolism

When *C. proridens* individuals are functioning as females, growth rate, as determined from length frequency analysis, slows due to metabolic changes (Dubovitsky 1977b). When fish change from female to male, growth rate increases. Fat content of the body cavity increases in mature fish to a maximum in November and December, just prior to spawning (Fig. 12) (Dubovitsky 1977a). Fat content decreases through the spawning season and is low by the end of spawning. Fat build-up is rapid in July and August in large post-spawning fish. During spawning, fat content is highest in fish living in deep water.

See 3.43.

3.5 Behavior

3.51 Migrations and local movements

No migrations of *C. arctifrons* have been reported.

Migrations of *C. proridens* on Campeche Bank have been reported by Dubovitsky (1977a). Eggs and larvae move primarily from east to west across the bank, and move towards shallow water (Fig. 13). Small fish move gradually from west to east in December through March. The majority of mature fish migrate into coastal waters of the central bank in January, dispersing eastward by March. Migration patterns in mature fish are strongly developed during the spawning period, with spawners generally moving offshore. After spawning, fish move into shallower water again. An extensive migration of 19-21 cm (length unspecified) fish to inshore waters occurs in January. Aggregations of *C. proridens* on Campeche Bank tend to form when bottom temperatures are lowest.

See 2.3 and 3.52.



Figure 13.—Schematic representation of migrations and movements of *Calamus proridens* on Campeche Bank. Arrows indicate movements of: 1, eggs and larvae; 2, immature individuals; 3, spawning individuals; 4, post-spawning individuals. (From Dubovitsky 1977a, fig. 10)



Figure 14.—Overall sex ratio of *Calamus proridens* in the commercially exploited population on Campeche Bank. (From Dubovitsky 1977b, fig. 2)

3.52 Schooling

Although schooling behavior is not well known in *C. arctifrons* and *C. proridens*, most small porgies form aggregations (Randall and Vergara R. 1978). Large individuals are more often solitary or found in small groups. Dubovitsky (1977a) noted concentrations of *C. proridens* on Campeche Bank, but stated that they did not form dense spawning aggregations.

See 3.51.

4 POPULATION

4.1 Structure

4.11 Sex ratio

Sex ratio of *C. arctifrons* is unknown.

On Campeche Bank, Dubovitsky (1977b) found that female *C. proridens* always predominated, regardless of time or location. Sex ratio observed was not constant from year to year, but averaged 1:2.7 males to females. About 73% of the commercially exploited population were females. Sex ratio changed drastically with size of the fish examined (Fig. 14). Only about 1% of specimens <23 cm (measurement unspecified) were males (Table 3). Beyond 24 cm, the percentage of males increased sharply, to about 69% of the total. Only females were found in the smallest specimens, and only males in the largest. Dubovitsky (1977b) derived the following linear relationship between sex ratio and size of fish:

$$S = 2.338 - 0.072\ L$$

where S is the proportion of females, and L is the mean length of the females examined. It is obvious from Fig. 14, however, that the true relationship approaches linearity only over a restricted range of length.

See 4.13.

4.13 Size composition

Size composition of *C. arctifrons* has not been reported.

Size composition of *C. proridens* on Campeche Bank was examined by Dubovitsky (1973, 1977a, b) (Figs. 15-20; Table 4). Size distributions presented by Dubovitsky (1977a) were often bimodal, with the smaller size-class dominating. In stocks of mature fish, females 18-22 cm (measurement unspecified) predominated (Fig. 20). About 59% of all females in the commercially exploited population were between 18 and 21 cm in length, and 65% of the males were between 25 and 28 cm. Size composition of females and their proportions in the commercial population are presented in Fig. 21.

See 2.3.

Table 3.—Percentage of females in three size ranges of *Calamus proridens* in the sexually mature part of the commercial catch on Campeche Bank, 1969-71. (From Dubovitsky 1977b)

Size range[1] (cm)	% females	% examined specimens
14-20	99.3	37.9
21-25	78.5	39.5
26-34	19.7	22.6

[1]Type of length measurement unspecified.



Figure 15.—Size composition of *Calamus proridens* on Campeche Bank (n = 665, $\bar{x}$ = 22.73 cm). Type of length measurement was unspecified. (From Dubovitsky 1973, fig. 3)



Figure 16.—Size composition of *Calamus proridens* on eastern Campeche Bank, by month. Solid lines represent specimens collected at depths <30-35 m, dashed lines represent specimens collected at >30-35 m. Cross-hatching represents specimens with ovaries in pre-spawning condition. Type of length measurement was unspecified. (From Dubovitsky 1977a, fig. 3)



Figure 17.—Size composition of *Calamus proridens* on central Campeche Bank, by month. Solid lines represent specimens collected at depths <30-35 m, dashed lines represent specimens collected at >30-35 m. Cross-hatching represents specimens with ovaries in pre-spawning condition. Type of length measurement was unspecified. (From Dubovitsky 1977a, fig. 3)



Figure 18.—Size composition of *Calamus proridens* on northwestern Campeche Bank, by month. Solid lines represent specimens collected at depths <30-35 m, dashed lines represent specimens collected at >30-35 m. Cross-hatching represents specimens with ovaries in pre-spawning condition. Type of length measurement was unspecified. (From Dubovitsky 1977a, fig. 4)



Figure 19.—Size composition of *Calamus proridens* on western Campeche Bank, by month. Solid lines represent specimens collected at depths <30-35 m, dashed lines represent specimens collected at >30-35 m. Cross-hatching represents specimens with ovaries in pre-spawning condition. Type of length measurement was unspecified. (From Dubovitsky 1977a, fig. 3)



Figure 20.—Size distribution of male and female *Calamus proridens* in the Campeche Bank commercial population with a histogram of lengths. (From Dubovitsky 1977b, fig. 1)



Figure 21.—Size distribution of female *Calamus proridens* and their frequency in the sexually mature part of the Campeche Bank population, 1969-1971. Type of length measurement was unspecified. (From Dubovitsky 1977b, fig. 3)

Table 4.—Size composition of *Calamus proridens* on Campeche Bank. Numbers above horizontal lines refer to specimens caught at depths ≤35 m; figures below lines refer to specimens caught at depths >35 m. (From Dubovitsky 1977a, Table 2)

Months	Size range[1] (cm)	Modal group (cm)	Rel. freq. of modal group (%)	Mean length (cm)	Percent length distribution in sample <20 cm	20-25 cm	>25 cm
Eastern Area							
Dec.	18-34	21-24	30	25.9	7	39	54
		27-31	40				
Jan.	18-32	22-23	24	25.0	3	57	40
		29-31	24				
Feb.-Mar.	13-32	20-23	58	22.9	8	70	22
Aug.		20-22	21				
	16-31	24-25	22	23.7	11	57	32
Nov.	18-30	19-22	55	22.0	25	54	21
Central Area							
Jan.	15-29	19-22	51	21.2	28	58	14
	16-26	18-19	51	18.6	67	30	3
Mar.	14-34			22.9	27	41	32
	15-29	19-20	43	20.8	42	47	11
June		20-21					
	16-29	25	14	22.3	9	79	12
		18-20	47				
July	16-30	24-26	29	21.5	37	53	10
	16-26	18	48	19.0	76	23	1
Nov.	15-33			23.6	18	47	35
	16-30	18-20	58	20.6	47	44	9
Northwest Area							
		22-24	42				
Dec.	19-29	26-27	26	24.6	1	59	40
	17-27	19-21	48	21.6	26	64	10
		24-26	27				
Jan.	13-30	21-23	45	22.4	15	65	20
July	15-27	20-22	55	21.6	15	77	8
		21-24	56				
Nov.	19-30	26-28	32	24.3	1	63	36
	15-30	18-21	56	20.8	40	51	9
Southwest Area							
Dec.	17-29	19-21	56	20.7	34	63	3
Jan.	15-34	22-23	35	24.5	3	66	31
July	17-26	20-21	63	21.0	11	88	1
	14-26	16-19	72	18.2	78	21	1

[1]Type of length measurement unspecified.



Figure 22.—Index of relative abundance of *Calamus arctifrons* on the West Florida Shelf, January 1978, expressed in terms of biomass and number of individuals. (From Darcy and Gutherz, unpubl., fig. 12.)

4.14 Subpopulations

No subpopulations of *C. arctifrons* or *C. proridens* are known.

4.2 Abundance and density

Calamus arctifrons—*Calamus arctifrons* was reported very abundant around the Florida Keys by Jordan and Fesler (1893). Tabb and Manning (1961) found *C. arctifrons* abundant on *Thalassia* flats in Florida Bay; they were most common from October to April. A trawling survey of the West Florida Shelf (Darcy and Gutherz 1984) collected *C. arctifrons* at 7.7% of the stations sampled; this species ranked tenth in biomass of all demersal fish sampled in 9-93 m in the survey area, and accounted for 3.1% of the total demersal fish biomass collected in that region. North of Tampa Bay, *C. arctifrons* ranked eighth in biomass in 9-35 m, and accounted for 5.6% of the total biomass. Abundance was highest in shallow (9-18 m) water (Fig. 22) and on rock bottom (Darcy and Gutherz unpubl.). A plot of mean abundance by grid zone on the West Florida Shelf (Fig. 23) shows maximum abundance in nearshore waters off the northwest coast of Florida. Estimated standing stock of *C. proridens* on the West Florida Shelf at depths >9 m is about 28,000 mt (Darcy and Gutherz unpubl.). In shallower water in the eastern Gulf of Mexico, *C. arctifrons* was reported common by Reid (1954), Grimes (1971), Moe and Martin (1965), Tabb and Manning (1961), and Grimes and Mountain (1971). The species is not common in the West Indies (Beebe and Tee-Van 1928; Randall and Caldwell 1966).

See 2.1 and 2.3.

Calamus proridens—*Calamus proridens* abundance on the Atlantic coast of the United States is poorly known, but Struhsaker (1969) reported it rather rare in the South Atlantic Bight.

In the Gulf of Mexico, *C. proridens* is not very common in the extreme northern gulf (Walls 1975), but is quite common on the northern central West Florida Shelf (Darcy and Gutherz unpubl.) (Fig. 24). Smith (1976) reported that *C. proridens* is the most common sparid at depths >30 m in the eastern Gulf of Mexico, and Smith et al. (1975) reported it as occasional on the Florida Middle Ground. *Calamus proridens* occurred at 9.2% of the stations sampled on the West Florida Shelf by Darcy and Gutherz (1984), and ranked eighth in biomass (3.7% of total demersal fish biomass) in 9-93 m over the entire survey area. It ranked seventh in biomass (6.5% of total demersal fish biomass) north of Tampa Bay in 9-35 m depths. Abundance was greatest in moderate (19-55 m) depths (Fig. 25) (Darcy and Gutherz unpubl.); estimated standing stock on the West Florida Shelf was about 34,000 mt.

On Campeche Bank, *C. proridens* is the most abundant sparid (Dubovitsky 1973, 1977b); along with *C. nodosus* and *C. bajonado*, *C. proridens* makes up the bulk of the commercial trawl fish catch on Campeche Bank. It has been estimated to comprise 10% of the benthic fish fauna of the bank (Sauskan and Olaechea 1974), and is second only to the tomtate, *Haemulon aurolineatum*, in abundance (Olaechea and Sauskan 1974; Sauskan and Olaechea 1974). According to Dubovitsky (1977a), *C. proridens* is the most important commercial fish on the bank. Catch rates of >500 kg/h trawling have been reported (Dubovitsky 1977a), but rates of <100 kg/h are more usual (Sauskan and Olaechea 1974) (Table 5). Estimated density on Campeche Bank by season, expressed in kilograms per hectare, was:

Winter	*Spring*	*Summer*	*Fall*	$\bar{x}$
1.79	5.50	5.65	1.74	3.74



Figure 23.—Mean catch rate of *Calamus arctifrons* by 20′ × 20′ grid zones on the West Florida Shelf, January 1978. Catches are expressed in mean numbers of individuals caught per 10-min tow. (From Darcy and Gutherz, unpubl., fig. 13)



Figure 24.—Mean catch rate of *Calamus proridens* by 20′ × 20′ grid zones on the West Florida Shelf, January 1978. Catches are expressed in mean numbers of individuals caught per 10-min tow. (From Darcy and Gutherz, unpubl., fig. 15)

Table 5.—Mean catch rates of *Calamus proridens* per hour of trawling of SRT-M trawlers on Campeche Bank, and percentage of the total fish catch in each part of the bank accounted for by *C. proridens*. (From Sauskan and Olaechea 1974.)

	Area of Campeche Bank					
	West		Central		East	
Depth (m)	Mean catch (kg/h)	% of total fish	Mean catch (kg/h)	% of total fish	Mean catch (kg/h)	% of total fish
15-30	45.5	12	95.0	14	69.9	6
30-50	53.5	15	54.0	8	27.1	6
50-120	0.0	0	13.9	5	60.9	14



Figure 25.—Index of relative abundance of *Calamus proridens* on the West Florida Shelf, January 1978, expressed in terms of biomass and numbers of individuals. (From Darcy and Gutherz, unpubl., fig. 14)

Table 6.—Estimated standing stock of porgies on Campeche Bank, based on Cuban trawl surveys. Figures for 1974-75 represent two different estimates. (From Klima 1976)

	Dates of surveys			
	1964-72	1974-75	1974-75	Mean
Standing stock (mt)	209,000	182,000	461,000	325,000

(Olaechea and Sauskan 1974). Sauskan and Olaechea (1974) estimated total mean abundance of *C. proridens* on Campeche Bank at 88,000 mt. Estimates of standing stock of all porgies on Campeche Bank are presented in Table 6.

Calamus proridens does not seem to be very common in the West Indies (Randall and Caldwell 1966), though Beebe and Tee-Van (1928) reported them quite common in Haitian fish markets.

See 2.1, 2.21, 2.22, 2.3, 3.16, 3.51, and 4.13.

4.3 Natality and recruitment

4.31 Reproduction rates

Reproduction rates of *C. arctifrons* and *C. proridens* are unknown, but *C. proridens* is a sequential spawner based on ovarian examination and histology (Dubovitsky 1977a) and may spawn more than once each spawning period.

See 3.15 and 3.16.

4.32 Factors affecting reproduction

Although there are no reports of factors affecting reproduction in *C. arctifrons*, temperature probably determines spawning season. Spawning in *C. proridens* is known to be temperature-dependent and to coincide with maximum plankton productivity (Dubovitsky 1977a).

See 3.16.

4.33 Recruitment

Little is known of recruitment in *C. arctifrons*. Wang and Raney (1971) collected a small specimen (26 mm SL) in March in Charlotte Harbor estuary, Florida.

Calamus proridens as small as 18 mm SL were collected on shallow grassflats off Lower Matecumbe Key, Florida, in April, and specimens as small as 21 mm SL were collected in March (Springer and McErlean 1962). On Campeche Bank, *C. proridens* are recruited to the commercial stock between ages I and II, and at lengths of about 14 cm (measurement unspecified) (Dubovitsky 1977a, 1977b).

See 2.21, 2.3, and 3.51

4.6 The population in the community and the ecosystem

Calamus arctifrons is a common member of grassbed communities in shallow water, and live-bottom communities in somewhat deeper water. Associates of *C. arctifrons* in shallow water probably include: pinfish, *Lagodon rhomboides*; spottail pinfish, *Diplodus holbrooki*; sheepshead, *Archosargus probatocephalus*; pigfish, *Orthopristis chrysoptera*; toadfish, *Opsanus* spp., seahorses and pipefishes, Syngnathidae; mojarras, Gerreidae; seatrouts, *Cynoscion* spp.; gobies, Gobiidae; triggerfishes and filefishes, Balistidae; puffers, Tetraodontidae; and porcupinefishes, Diodontidae. In deeper water (>9 m), cluster analysis of demersal fish catches on the West Florida Shelf (Darcy and Gutherz unpubl.) grouped conger eels, *Hildebrandia* sp.; spottail pinfish; crested cusk-eel, *Ophidion welshi*; sand perch, *Diplectrum formosum*; scrawled cowfish, *Lactophrys quadricornis*; black sea bass, *Centropristis striata*; and shortnose batfish, *Ogcocephalus nasutus*, with *C. arctifrons*.

Calamus proridens is a member of live-bottom communities in shallow to moderate depths. On the West Florida Shelf, cluster analysis grouped tomtate; gulf flounder, *Paralichthys albigutta*; Atlantic croaker, *Micropogonias undulatus*; and the cusk-eel, *Lepophidium brevibarbe*, with *C. proridens* in water >9 m deep (Darcy and Gutherz unpubl.). On Campeche Bank, *C. proridens* is often caught with *C. penna*, *C. nodosus*, and *C. bajonado*, especially in summer (Olaechea and Sauskan 1974).

See 3.3.

5 EXPLOITATION

5.1 Fishing equipment

Calamus arctifrons is caught in beach seines (Evermann and Marsh 1902; Randall and Caldwell 1966) and bottom trawls, and on hook-and-line and bottom longlines (Randall and Vergara R. 1978). *Calamus proridens* is caught in fish traps, trawls, and on hook-and-line (Randall and Vergara R. 1978).

5.2 Fishing areas

Calamus arctifrons is fished in coastal waters of the Gulf of Mexico from southern Florida to Louisiana (Randall and Vergara R. 1978), but there is no directed fishery.

Calamus proridens is fished in coastal waters of Florida, Campeche Bay, and Cuba (Randall and Vergara R. 1978). Concentrations of *C. proridens* on Campeche Bank in summer were mapped by Olaechea and Sauskan (1974) (Fig. 26).

See 2.1, 2.22, 2.3, and 3.51.



Figure 26.—Principal zones of concentration of *Calamus proridens* on Campeche Bank in summer. *Calamus bajonado*, *C. penna* and *C. nodosus* are also concentrated in the same areas. (From Olaechea and Sauskan 1974, fig. 3)

5.3 Fishing seasons

No specific information regarding fishing seasons is available. Since there is no directed fishery for *C. arctifrons* or *C. proridens* in the United States, catches of these species are probably dependent upon effort expended in pursuit of other species. On Campeche Bank, *C. proridens* are present year-round, but are most concentrated, and therefore most easily fished, when bottom temperatures are low (Dubovitsky 1977a).

See 2.3 and 4.13.

5.4 Fishing operations and results

5.43 Catches

Calamus arctifrons and *C. proridens* are good food fish and are marketed mostly fresh and frozen (Evermann and Marsh 1902; Randall and Vergara R. 1978). Moe (1963) stated that these species are of minor commercial importance, but are sometimes caught by commercial snapper fishermen off the west coast of Florida, and are often used as fresh bait. *Calamus proridens* is one of the most important commercial species on Campeche Bank (Sokolova 1965), and is not uncommon in markets in Haiti (Beebe and Tee-Van 1928).

Catch statistics for *C. arctifrons* and *C. proridens* are not reported separately (Randall and Vergara R. 1978). Total reported catch of *Calamus* spp. from FAO fishing area 31 (western central Atlantic Ocean) in 1975 was about 6,777 mt, but actual landings were undoubtedly much higher, since landings of porgies are usually combined with landings of other percomorph fishes (Randall and Vergara R. 1978). Catches of all species of porgies combined in FAO fishing area 31 were: 1970, 1,400 mt; 1971, 3,600 mt; 1972, 8,900 mt; 1973, 3,600 mt (Klima 1976). Catches of porgies by individual nations fishing in the western central Atlantic area in 1970-73 are shown in Table 7. Most of the reported catch was probably from Campeche Bank. Porgies, of which *C. proridens* is the most common, comprised 30% of trawl survey catches by the Cubans on Campeche Bank in the early 1970's (Klima 1976). Based on these and other surveys, Campeche Bank could yield 33,000-65,000 mt of porgies each year (Klima 1976). Sauskan and Olaechea (1974) estimated the potential catch of *C. proridens* on the bank as 15,000-20,000 mt. Porgies made up 56-58% of Cuban trawl catches on Campeche Bank in 1973-74 (Klima 1976). Commercial catches on the bank have been quite variable, possibly due to changes in environmental conditions (Klima 1976).

Recreational catch figures, like commercial catch figures, are seldom reported for separate species of porgies. The total recreational catch of porgies in the Gulf of Mexico in 1960 was estimated at 8,550,000 fish, of 5,792,000 kg (Clark 1962). Most (72%) were caught from boats, and the rest from shore. Recreational catches of porgies reported in 1970 were 13,234,000 fish, of 9,671,000 kg, in the eastern Gulf of Mexico, and 16,230,000 fish, of 10,913,000 kg, along the Atlantic coast of the United States south of Cape Hatteras (Deuel 1973). Landings are stratified by fishing area and method in Table 8.

See 2.3 and 4.2

Table 7.—Reported catches of porgies from the western central Atlantic fishing area, 1970-73, by fishing nation. (From Klima 1976)

	Catch per year (mt)			
Nation	1970	1971	1972	1973
Cuba	400	2,000	1,900	1,900
U.S.S.R.	0	400	6,000	300
Mexico	500	600	600	800

Table 8.—Recreational catches of porgies in the eastern Gulf of Mexico and along the southern Atlantic coast of the United States, 1970. Figures are in thousands of individuals caught. (Data from Deuel 1973)

	Fishing area		Fishing method			
Geographical area	Ocean	Sounds, rivers, bays	Private, rented boat	Party, charter boat	Bridge, pier, jetty	Beach, bank
South Atlantic	6,266	9,964	4,989	1,068	7,499	2,674
Eastern Gulf of Mexico	905	12,329	6,579	128	4,587	1,940

6 PROTECTION AND MANAGEMENT

6.1 Regulatory measures

Calamus arctifrons is included in the Snapper/Grouper Complex Fishery Management Plan of the South Atlantic Fishery Management Council. Both *C. arctifrons* and *C. proridens* are included in the Reef Fish Fishery Management Plan of the Gulf of Mexico Fishery Management Council.

ACKNOWLEDGMENTS

I thank Donald M. Allen and Albert C. Jones of the National Marine Fisheries Service, Southeast Fisheries Center, for critically reviewing this manuscript.

CITATIONS

BEEBE, W., and J. TEE-VAN.
1928. The fishes of Port-au-Prince Bay, Haiti, with a summary of the known species of marine fish of the island of Haiti and Santo Domingo. Zoologica (N.Y.) 10:1-279.

BREDER, C. M., Jr.
1962. On the significance of transparency in osteichthid fish eggs and larvae. Copeia 1962:561-567.

BRIGGS, J. C.
1958. A list of Florida fishes and their distribution. Bull. Fla. State Mus. Biol. Sci. 2:223-318.

CLARK, J. R.
1962. The 1960 salt-water angling survey. U.S. Fish Wildl. Serv. Circ. 153, 36 p.

CUVIER, G., and A. VALENCIENNES.
1830. Histoire naturelle des poissons. F. G. Levrault, Paris, vol. 6, 559 p. (Not seen)

DARCY, G. H., and E. J. GUTHERZ.
Unpubl. Diversity, distribution, abundance, and communities of demersal fishes and selected invertebrates on the West Florida Shelf, January 1978. Southeast Fisheries Center, National Marine Fisheries Service, NOAA, 75 Virginia Beach Drive, Miami, FL 33149-1099, 327 p.
1984. Abundance and density of demersal fishes on the West Florida Shelf, January 1978. Bull. Mar. Sci. 34:81-105.

DEUEL, D. G.
1973. 1970 salt-water angling survey. U.S. Dep. Commer. Current Fish. Stat. 6200, 54 p.

DUARTE-BELLO, P. P.
1959. Catalogo de peces Cubanos. Univ. de Villanueva (Cuba). Monogr. 6:1-208.

DUBOVITSKY, A. A.
1973. The specific composition of sparids, the genus *Calamus*, family Sparidae, on the Campeche Bank. [In Russ.] Tr. Vses. Nauchno-Issled. Inst. Morsk. Ryb. Khoz. Okeanogr. 93:230-244.
1977a. Distribution, migrations and some biological features of littlehead porgy (*Calamus proridens* Jordan and Gilbert, 1884) family Sparidae, of the Gulf of Mexico. *In* H. B. Stewart (editor), Cooperative investigations of the Caribbean and adjacent regions — II, Caracas, Venezuela, 12-16 July 1976, p. 123-143. FAO Fish Rep. 200.
1977b. Sex ratio and length-sex composition of the Campeche-Bank littlehead porgy (*Calamus proridens*, Jordan and Gilbert, 1884) population. *In* H. B. Stewart (editor), Cooperative investigations of the Caribbean and adjacent regions — II, Caracas, Venezuela, 12-16 July 1976, p. 115-122. FAO Fish Rep. 200.

EVERMANN, B. W., and M. C. MARSH.
1902. The fishes of Puerto Rico. Bull. U.S. Fish. Comm. 20, Pt. 1, 1900: 49-350.

GONZÁLES, O.C.
1973. Dependencia afloramiento - capturas y pronóstico a corta plaza en el región oriental del Banco de Campeche. Inst. Nac. Pesca, Cuba, Inf. Invest. 6:1-6 (Not seen)

GOODE, G. B., and T. H. BEAN.
1882. Descriptions of twenty-five new species of fish from the southern United States, and three new genera, *Letharcus*, *Ioglossus*, and *Chriodorus*. Proc. U.S. Nat. Mus. 5:412-437.

GREGORY, W. K.
1933. Fish skulls, a study of the evolution of natural mechanisms. Trans. Am. Philosoph. Soc. 23:75-481.

GRIMES, C. B.
1971. Thermal additional studies of the Crystal River steam electric station. Fla. Dep. Nat. Resour. Mar. Lab., Prof. Pap. Ser. 11, 53 p.

GRIMES, C. B., and J. A. MOUNTAIN.
1971. Effects of thermal effluent upon marine fishes near the Crystal River steam electric station. Fla. Dep. Nat. Resour. Mar. Res. Lab., Prof. Pap. Ser. 17, 64 p.

GUICHENOT, A.
1868. Révision du genre des pagels (*Pagellus*, *Lithognathus*, *Calamus*). Mém. Soc. Impér. Sci. Natur. Cherbourg 14:97-123. (Not seen)

HASTINGS, R. W., L. H. OGREN, and M. T. MABRY.
1976. Observations on the fish fauna associated with offshore platforms in the northeastern Gulf of Mexico. Fish. Bull., U.S. 74:387-402.

HOESE, H. D., and R. H. MOORE.
1977. Fishes of the Gulf of Mexico, Texas, Louisiana, and adjacent waters. Tex. A&M Univ. Press, College Station, 327 p.

HOUDE, E. D., J. C. LEAK, C. E. DOWD, S. A. BERKELEY, and W. J. RICHARDS.
1979. Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. Rep. to Bur. Land Manage. under contract no. AA550-CT7-28, June 1979, 546 p.

JORDAN, D. S., and B. W. EVERMANN.
1896-1900. The fishes of North and Middle America. Bull. U.S. Nat. Mus. 47:1-3313.

JORDAN, D. S., and B. FESLER.
1893. A review of the sparoid fishes of America and Europe. Rep. Comm., U.S. Comm. Fish Fish., Pt. 17, 1889-1891:421-544.

JORDAN, D. S., and C. H. GILBERT.
1882. Synopsis of the fishes of North America. Bull. U.S. Nat. Mus. 16:1-1018.
1884. A review of the species of the genus *Calamus*. Proc. U.S. Nat. Mus. 7:14-24.

JRÓMOV, N. S.
1967. Sobre la investigación del plancton en èl Golfo de México y el Mar Caribe. VNIRO/CIP Invest. Pesq. Soviético-Cubanas 2:39-57. (Not seen)

JUÁREZ, M.
1975. Distribucion cuantitativa y algunas aspectos cualitativos del ictioplancton del Banco de Campeche. Rev. Invest. Inst. Nac. Pesca (Cuba) 1(1):27-71.

KLIMA, E. F.
1976. An assessment of the fish stocks and fisheries of the Campeche Bank. WECAF Stud. 5, FAO, Rome, 24 p.

LONGLEY, W. H., and S. F. HILDEBRAND.
1941. Systematic catalogue of the fishes of Tortugas, Florida, with observations on color, habits, and local distribution. Carnegie Inst. Wash. Publ. 535:1-331.

MILLER, G. C., and W. J. RICHARDS.
1980. Reef fish habitat, faunal assemblages and factors determining distributions in the South Atlantic Bight. Proc. Gulf Caribb. Fish. Inst., 32nd Annu. Meet., p. 114-130.

MOE, M. A.
1963. A survey of offshore fishing in Florida. Fla. Board Conserv. Mar. Res. Lab. Prof. Pap. Ser. 4, 117 p.

MOE, M. A., Jr., and G. T. MARTIN.
1965. Fishes taken in monthly trawl samples offshore of Pinellas County, Florida, with new additions to the fish fauna of the Tampa Bay area. Tulane Stud. Zool. 12:129-151.

NAHHAS, F. M., and R. M. CABLE.
1964. Digenetic and aspidogastrid trematodes from marine fishes of Curaçao and Jamaica. Tulane Stud. Zool. 11:169-228.

OLAECHEA, A.
1975. Distribucion y abundancia de peces demersales y su relacion con la temperatura de las aguas del fondo. Resum. Invest. Inst. Nac. Pesca Cent. Invest. Pesq., Cuba 2:161-164.

OLAECHEA, A., and V. I. SAUSKAN.
1974. Cartas de pesca del Banco de Campeche, 1972. Resum. Invest. Inst. Nac. Pesca Cent. Invest. Pesq., Cuba 1:176-181.

POEY, F.
1874. Monographie des poissons de Cuba compris dans la sousfamille des Sparini. Ann. Lyc. Nat. Hist. N.Y. 10:170-184. (Not seen)

RANDALL, J. E., and D. K. CALDWELL.
1966. A review of the sparid fish genus *Calamus*, with descriptions of four new species. Bull. Los Angeles Co. Mus. Nat. Hist., Sci. 2:1-47.

RANDALL, J. E., and R. VERGARA R.
1978. Sparidae. *In* W. Fischer (editor), FAO species indentification sheets for fishery purposes, Western Central Atlantic (Fishing Area 31), Vol. 4, unpaginated. FAO. Rome.

REID, G. K., Jr.
1954. An ecological study of the Gulf of Mexico fishes, in the vicinity of Cedar Key, Florida. Bull. Mar. Sci. Gulf Caribb. 4:1-94.

ROBINS, C. R., R. M. BAILEY, C. E. BOND, J. R. BROOKER, E. A. LACHNER, R. N. LEA, and W. B. SCOTT.
1980. A list of common and scientific names of fishes from the United States and Canada. 4th ed. Am. Fish. Soc. Spec. Publ. 12, 174 p.

de la SAGRA, R.
1843. Historia física, política y natural de la isla de Cuba. Arthus Bertrand, Paris, 4:145-254. (Not seen)

SAUSKAN, V. I., and A. OLAECHEA.
1974. Ictiofauna bentonica del Banco de Campeche. Resum. Invest. Inst. Nac. Pesca Cent. Invest. Pesq., Cuba 1:102-106.

SMITH, G. B.
1976. Ecology and distribution of eastern Gulf of Mexico reef fishes. Fla. Mar. Res. Publ. 19, 78 p.

SMITH, G. B., H. M. AUSTIN, S. A. BORTONE, R. W. HASTINGS, and L. H. OGREN.
1975. Fishes of the Florida Middle Ground with comments on ecology and zoogeography. Fla. Mar. Res. Publ. 9, 14 p.
SMITH, G. B., D. A. HENSLEY, and H. H. MATHEWS.
1979. Comparative efficacy of artificial and natural Gulf of Mexico reefs as fish attractants. Fla. Mar. Res. Publ. 35, 7 p.
SOKOLOVA, L. V.
1965. Distribution and biological characteristics of the main commercial fish of Campeche Bank. *In* A. S. Bogdanov (editor), Soviet-Cuban fishery investigations, p. 208-224. (Engl. transl. by Israel Prog. Sci. Transl., Jerusalem, 1969.)
SPRINGER, V. G., and A. J. McERLEAN.
1962. Seasonality of fishes on a south Florida shore. Bull. Mar. Sci. Gulf Caribb. 12:39-60.
SPRINGER, V. G., and K. D. WOODBURN.
1960. An ecological study of the fishes of the Tampa Bay area. Fla. Dep. Nat. Resour. Mar. Res. Lab., Prof. Pap. Ser. 1, 104 p.
STRUHSAKER, P.
1969. Demersal fish resources: composition, distribution, and commercial potential of the continental shelf stocks off southeastern United States. U.S. Fish Wildl. Serv., Fish. Ind. Res. 4:261-300.
TABB, D. C., and R. B. MANNING.
1961. A checklist of the flora and fauna of northern Florida Bay and adjacent waters of the Florida mainland collected during the period of July, 1957 through September, 1960. Bull. Mar. Sci. Gulf Caribb. 11:552-649.
VICK, N. G.
1964. The marine ichthyofauna of St. Andrew Bay, Florida, and nearshore habitats of the northeastern Gulf of Mexico. Tex. A&M Univ. Res. Found., A&M Proj. 286-D, 77 p.
WALLS, J. G.
1975. Fishes of the northern Gulf of Mexico. T.F.H. Publ., Inc., Neptune City, N.J., 432 p.
WANG, J. C. S., and E. C. RANEY.
1971. Distribution and fluctuations in the fish fauna of the Charlotte Harbor estuary, Florida. Charlotte Harbor Estuarine Study, Mote Mar. Lab., 56 p.
ZANEVELD, J. S.
1983. Caribbean fish life. E. J. Brill, Leiden, 163 p.

NOAA TECHNICAL REPORT NMFS

Guidelines for Contributors

FORM OF MANUSCRIPT

Authors are encouraged to retain manuscripts on word processing storage media, such as diskettes, floppy disks, cassette or magnetic tapes, and submit a hard copy run from the storage media. Hardcopy should be typed double-spaced on white bond paper in the following sequence:

Title page
Contents
Abstract
Text
Acknowledgments
Citations
Text footnotes
Appendix
Tables
Figure Legends
Figures

CONTENTS OF MANUSCRIPT

Contents. Contains all text headings with page numbers omitted.

Abstract. Not to exceed one double-spaced typed page. Should include a sentence or two explaining to the general reader why the research was undertaken and why the results should be viewed as important. Abstract should convey the main point of the paper and outline the results or conclusions. Footnotes and references do not belong in the abstract.

Text. A brief introduction should portray the broad significance of the paper. The entire text should be intelligible to readers from different disciplines. All technical terms should be defined. Follow the *U.S. Government Printing Office Style Manual*, 1984 edition.

Fishery nomenclature follows the American Fisheries Society Special Publication No. 12, *A List of Common and Scientific Names of Fishes from the United States and Canada*, 4th edition, 1980.

Abbreviations and symbols. Define all symbols, abbreviations, and acronyms. Define all symbols in equations and formulas. Abbreviate units of measure only when used with numerals or in tables and figures where there is lack of space. Periods are rarely used except for et al., vs., e.g., i.e., etc.

Measurements should be expressed in metric units. Other equivalent units may be given in parentheses.

Text footnotes. Type on a separate sheet from the text. Footnotes are not used for reference material or personal communications, but rather to explain or define terms in the text and for contribution numbers.

Personal communications are noted in parentheses in the text (name, affiliation, brief address including zip code, month and year).

Citations. All written sources should be listed in the Citations section, including unpublished and processed material. In text, cite as Smith and Jones (1977) or (Smith and Jones 1977); if more than one citation, list chronologically (Smith 1936; Jones 1975; Doe 1977). All sources cited in the text should be listed alphabetically by the senior authors' surname under the heading CITATIONS. Abbreviations of periodicals and serials should conform to the Biosis Data Base™. The author is responsible for the accuracy of all citations.

Acknowledgments. Gather all acknowledgments into a brief statement at the end of the text. Give credit only for exceptional contributions and not to those whose contributions are part of their normal duties.

Tables. Tables should supplement, not duplicate, the text. Each table should appear on a separate page, and be numbered consecutively. Headings should be short but amply descriptive so that the reader need not refer to the text. For values less than 1, zeros should precede all decimal points. Give each column a heading with units of measure indicated in parentheses. Do not change the unit of measure within a column. Table footnotes should be numbered consecutively in Arabic numerals across the page from left to right and down; to avoid confusion with exponents, place them to the left of numerals, e.g., $^{3}10$. All tables should be cited consecutively in the text and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page. Lengthy tables that are typed in the proper format and are clean and legible can be used camera-ready.

Figures. Photographs and line drawings should be of professional quality—clear and concise—and reducible to 42 picas for full-page width or to 20 picas for a single-column width, and a maximum of 57 picas high. All illustrations must be proportioned so that they are legible when reduced to fit the page format. Line weight and lettering should be sharp and even. Lettering on graphs should be upper and lower case, and vertical lettering should be avoided as much as possible (except for vertical, y, axis). For values less than 1, zeros should precede all decimal points. Reproductions of line art work are accepted in the form of photographic prints from negatives or photomechanical transfer. Halftones should be sharply focused with good contrast. Micron rules should be inserted on electron micrographs, even when magnification is included in the figure legend. There should be good distinction between identifying letters (preferably press-on) and background of photograph. All figures should be cited consecutively in the text and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page. Each figure should be labeled in pencil on the back. DO NOT SEND original or photographic prints of figures to the Scientific Editor; they will be requested when the manuscript is accepted for publication.

Copyright. Government publications are in the public domain, i.e., they are not protected by copyright.

SUBMISSION OF MANUSCRIPTS

Send original hardcopy and two duplicated copies of the manuscript to:

Dr. William J. Richards, Scientific Editor
NOAA Technical Reports NMFS
Southeast Fisheries Center
National Marine Fisheries Service, NOAA
75 Virginia Beach Drive
Miami, FL 33149

Indicate whether word processing storage media are available, but do not forward disks until requested by Scientific Editor or Scientific Publications office.

Copies of published reports. For single monographs, the senior author will be provided with 50 copies free of charge and 50 to his or her organization. For collections of papers, each senior author will receive 50 separates and one cover, and his or her organization will receive the same. Additional copies may be purchased in lots of 100.

ADDITIONAL INFORMATION

Questions regarding editorial policy and procedures may be forwarded to:

Nancy Peacock, Managing Editor
NOAA Technical Reports NMFS
National Marine Fisheries Service, NOAA
Scientific Publications Office
Bin C15700
Seattle, WA 98115

FISHERIES SYNOPSES

This series of documents, issued by FAO, CSIRO, INP, and NMFS, contains comprehensive reviews of present knowledge on species and stocks of aquatic organisms of present or potential economic interest. The Fishery Resources and Environment Division of FAO is responsible for the overall coordination of the series. The primary purpose of this series is to make existing information readily available to fishery scientists according to a standard pattern, and by so doing also to draw attention to gaps in knowledge. It is hoped that synopses in this series will be useful to other scientists initiating investigations of the species concerned or of related ones, as a means of exchange of knowledge among those already working on the species, and as the basis for comparative study of fisheries resources. They will be brought up to date from time to time as further information becomes available.

The documents of this series are issued under the following titles:

		Symbol
FAO	**Fisheries Synopsis No.**	**FIR/S**
CSIRO	**Fisheries Synopsis No.**	**DFO/S**
INP	**Sinopsis sobre la Pesca No.**	**INP/S**
NMFS	**Fisheries Synopsis No.**	**NMFS/S**

Synopses in this series are compiled according to a standard outline described in FIb/S1 Rev. 1 (1965). FAO, CSIRO, INP, and NMFS are working to secure the cooperation of other organizations and of individual scientists in drafting synopses on species about which they have knowledge, and welcome offers of help in this task. Additions and corrections to synopses already issued will also be most welcome. Comments on individual synopses and requests for information should be addressed to the coordinators and editors of the issuing organizations, and suggestions regarding the expansion or modification of the outline to FAO:

FAO:

Fishery Resources and Environment Division
Aquatic Resources Survey and Evaluation Service
Food and Agriculture Organization of the United Nations
Via delle Terme di Caracalla
00100 Rome, Italy

CSIRO:

CSIRO Division of Fisheries and Oceanography
Box 21
Cronulla, N.S.W. 2230
Australia

INP:

Instituto Nacional de Pesca
Subsecretaria de Pesca
Secretaria de Pesca
Secretaria de Industria y Comercio
Carmona Y Valle 101-403
Mexico 7, D.F.

NMFS:

Scientific Editor
Southwest Fisheries Center
National Marine Fisheries Service, NOAA
P.O. Box 271
La Jolla, CA 92038
U.S.A.

Consolidated lists of species or groups covered by synopses issued to date or in preparation will be issued from time to time. Requests for copies of synopses should be addressed to the issuing organization; except for NMFS/S copies, these can be purchased from National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22151.

The following synopses in this series have been issued since January 1983:

NMFS/S 133	Synopsis of biological data on the grunts *Haemulon aurolineatum* and *H. plumieri* (Pisces: Haemulidae)	February 1983
NMFS/S 134	Synopsis of biological data on the pigfish, *Orthopristis chrysoptera* (Pisces: Haemulidae)	March 1983
NMFS/S 136	Synopsis of biological data on skipjack tuna, *Katsuwonus pelamis*	January 1984
NMFS/S 138	Synopsis of biological data on the blue crab, *Callinectes sapidus* Rathbun	March 1984
NMFS/S 140	Synopsis of biological data on shortnose sturgeon, *Acipenser brevirostrum* LeSueur 1818	October 1984
NMFS/S 141	Synopsis of biological data on the pinfish, *Lagodon rhomboides* (Pisces: Sparidae)	February 1985
NMFS/S 142	Synopsis of biological data on the spottail pinfish, *Diplodus holbrooki* (Pisces: Sparidae)	January 1985
NMFS/S 143	Synopsis of biological data on the sand perch, *Diplectrum formosum* (Pisces: Serranidae)	March 1985
NMFS/S 144	Synopsis of biological data on the pink shrimp, *Pandalus borealis* Krøyer 1838	May 1985

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC
ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS OFFICE
BIN C15700
SEATTLE, WA 98115

OFFICIAL BUSINESS

Penalty for Private Use, $300

BULK RATE
POSTAGE & FEES PAID
U.S. Department of Commerce
Permit No. G-19

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 13, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS—Reports containing data, observations, instructions, etc. A partial listing includes data serials; predictions and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

Information on availability of NOAA publications can be obtained from:

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161